Vegetables guiding program

(Onion)

Pre-planting herbicide

To be applied before tilling the soil to control the weeds:

Use one product of the below:

Herbicide	Mode of action	Rate of use/ha	Dosage/15L
Force up liquid	Non selective systemic herbicide	4 L	200 ml
	systemic nerbicide		
Force up granule	Non selective	1,5kg	50 g
	systemic herbicide		
Parae Force	Non selective contact	4 L	200 ml
	herbicide		

Pre-emergence herbicide

Use one product of the below:

To be applied immediately after transplanting

Herbicide	Mode of action	Rate of use/ha	Dosage/15L
Force top	Selective pre emergence herbicides	3 L	160 ml
Albasa force	for vegetables	1 L	100 ml

Post-emergence Herbicide

To be applied three weeks after transplanting

Use the below to control the weeds

Herbicide	Mode of action	Rate of use/ha	Dosage/15L
Benta force	Selective post emergence herbicide for vegetables to control broad leaf weeds , yellow nut sedges	3 L	120 ml

Insecticides

Item	Mode of action	Rate of use/ha	Dosage/15L
	Contact insecticide		
Iron force	& larvicide used to control worms ,	500 ml -1L	20ml-40ml
	diamond back moth		
	Contact & systemic insecticide used to		
Laraforce gold	control worms, aphids, grass	500 ml	25ml
	hoppers, thrips, and white flies in		
	vegetables		

Fungicide

	vegetables		
Fungicide			
Item	Mode of action	Rate of use/ha	Dosage/15L
Green force	Preventive & curative fungicide used to control powdery mildew, downy mildew, leaf blight, leaf spot diseases, fusarium wilt and seed dressing.	500g	50g
Total force	Systemic fungicide used to control fungal diseases in vegetables	400g	20g

foliar fertilization program

Item	Mode of application	Rate of use/ha	Dosage/15L
T- maxi		250ml	25ml
C- maxi	Foliar application	250ml	25ml
Maxi K		2 kg	50 g
Boostract		500g	50 g

To be sprayed within 5-7 days interval between the applications

Surfactant shall be mixed with the foliar applied chemical to increase efficiency and to reduce the chemical wash off effect by rain

Surfactant

Item	Mode of action	Rate of use/ha	Dosage/15L
Cover force	Maximize spreading and wetting for better uptake of agrochemical and fertilizers by the crops leaves. Reduce the washing of chemical by rain. To be used mixed with all the items above except the pre-emergence herbicides	100 ml	5 ml