Cotton Guiding Program

Pre-planting herbicide

To be applied before planting the crop seeds to control the geminated weeds:

Use one product of the below:

Herbicide	Mode of action	Rate of use/ha	Dosage/15L tecto sprayer
Force up liquid	Non- selective systemic herbicide	4 L	200 ml
Force up granule	Non- selective systemic herbicide	1.5 kg	50 g
United force	Non- selective systemic herbicide	4L	150 ml
Parae Force	Non- selective contact herbicide	41	200 ml

Pre-emergence herbicide

Use one product of the below:

To be applied after planting the crop seeds before the germination:

Herbicide	Mode of action	Rate of use/ha	Dosage/15L tecto sprayer
Trace force	Selective systemic pre- emergence herbicide for	2 L	150 ml
Force top	cotton.	2 L	150 ml
Force-uron		4 L	160 ml
Buta force		2 kg	100 g

Post-emergence Herbicide

To be applied in cotton farm to control the germinated weeds

Herbicide	Mode of action	Rate of use/ha	Dosage/15Ltecto
			sprayer
	Selective systemic post		
Star force	emergence herbicide for cotton to control	1 L	70 ml
	grasses.		

Insecticides

Item	Mode of action	Rate of use/ha	Dosage/15Ltecto
			sprayer
Punch	Contact insecticide to control spider mite	11	40 ml
Hill force	Contact insecticide to control termites, stink bugs, spider mites	250 ml	25 ml
Final force	Systemic insecticide used to control boll worm, stink bug	600 g	30 g

Fungicide

Item	Mode of action	Rate of use/ha	Dosage/15Ltecto
			sprayer
	Systemic Curative	400 g	20 g
Total force	fungicide against leaf spot disease.		

Fungicare	Preventive and curative	500 ml	30 ml
	fungicide against		
	anthracnose disease		

Foliar fertilization program

To be sprayer with 15 days interval between the applications

Item	Mode of application	Rate of use/ha	Dosage / 15 L tecto
			sprayer
Boostract	Foliar applied fertilizer used to increase productivity	11	50 ml

Surfactant to be mixed with the foliar applied chemical to increase efficiency

Surfactant

Item	Mode of action	Rate of use/ha	Dosage / 15 L tecto
			sprayer
Cover force	Maximize spreading and wetting for better uptake of agrochemical and fertilizers by the crops leaves. Reduce the washing of chemical by rain. To be used mixed with all the items above except the preemergence	100 ml	5 ml

